

United States Department of Agriculture National Agricultural Statistics Service

FLORIDA CROP PROGRESS & CONDITION REPORT



In cooperation with the Florida Department of Agriculture and Consumer Services and the UF/IFAS Extension Service 2290 Lucien Way, Suite 300, Maitland, FL 32751 · (407) 648-6013 · (855) 271-9801 FAX · www.nass.usda.gov/fl

Released: January 12, 2015 (4 PM EST) Week Ending: January 11, 2015

Cold Nights, Warm Days, Little Rain

Weather Summary: According to Florida's Automated Weather Network (FAWN), rainfall ranged from none to 0.84 of an inch across the State. Fort Lauderdale (Broward County) received the most rain with 0.84 of an inch. Per the U.S. Drought Monitor, Florida was 94 percent drought free this past week. Temperatures ranged from 17 degrees for night time lows to 85 degrees for day time highs, across the State. The highest temperature was 85 degrees in Immokalee (Collier County) and Fort Lauderdale (Broward County). The lowest temperature in the State was 17 degrees in Defuniak Springs (Walton County) and Jay (Santa Rosa County).


Soil Moisture Ratings

	Topsoil			
Moisture Rating	Current week	Previous week	Previous year	
	(percent)	(percent)	(percent)	
Very short	1	0	3	
Short	22	19	15	
Adequate	73	76	75	
Surplus	4	5	7	


Field Crops: There was an average of 6.4 days suitable for field work this past week, slightly above the previous week. Seasonal field preparation in the Panhandle continued this past week.

Fruit and Vegetables: Flagler, Putnam, and Bradford county farmers continued harvesting cabbage and cold weather crops. Potato planting continued in Flagler and Putnam counties. In southwest Florida, foggy mornings were causing disease problems in vegetables. Vegetables coming to market in southwest Florida were collards, cucumbers, eggplant, mustard, peppers, squash, tomatoes, and a variety of specialty items. Crops harvested in Miami-Dade County were green beans, pole beans, yellow squash, zucchini, tomatoes, peppers, eggplant, sweet corn, boniato, malanga, and avocado.

Precipitation (in)-Florida: Jan. 5-11, 2015



U.S. Drought Monitor-Florida: Jan. 6, 2015



Livestock and Pastures: In the Panhandle, cold weather hurt winter grazing. Cattlemen were feeding hay. In Charlotte, Collier, Glades, Hendry, and Lee counties, pasture quality was aided by warm temperatures but livestock producers provided supplemental feed to cattle as necessary. Statewide, the cattle condition was mostly good while the pasture condition was fair to good.

Cattle and Pasture Condition

	Cattle		Pasture	
Condition	Current week	Previous week	Current week	Previous week
	(percent)	(percent)	(percent)	(percent)
Very poor	0	0	3	2
Poor	3	3	13	14
Fair	29	26	40	39
Good	65	67	44	45
Excellent	3	4	0	0

Citrus: Temperatures were cooler this past week in citrus producing counties. Highs ranged from the mid 70s to low 80s, and lows were in the 30s and 40s much of the week. Rainfall was very light. Seven of nineteen monitored stations did not record any rainfall. Only two stations recorded more than a tenth of an inch of rainfall, Umatilla (Lake County) at 0.14 of an inch and Immokalee (Collier County) at 0.15 of an inch. As per the U.S. Drought Monitor, last updated January 6, 2015, no drought or abnormally dry conditions exist in any citrus producing area.

Processing plants were up and running at full capacity. Harvesting for early and midseason oranges included both Hamlin and Pineapple oranges. Weekly totals were slightly less than last season, due to the small size fruit and lagging maturity levels.

Navel orange harvest was slowing as the season was coming closer to an end. Colored and white grapefruit harvest was overall running at a good pace and will continue for several more weeks. Honey tangerine harvest had started but was being harvested in small amounts.

Grove activity included spraying, irrigation on several days during the month, and mowing in preparation for harvest. Combating greening is a constant struggle for Citrus growers. Re-entry times after spraying is between two and ten days, slowing down survey progress at times.

Citrus Estimated Boxes Harvested

[In thousands of 1-3/5 bushel boxes]

	For week ending:			
Сгор	Dec 28, 2014	Jan 4, 2015	Jan 11, 2015	
	(boxes)	(boxes)	(boxes)	
Early & mid oranges	1.960	3,660	4,010	
Temples	0	0	3	
Navel oranges	19	23	28	
White grapefruit	20	41	45	
Colored grapefruit	161	268	374	
Sunburst tangerines	30	42	30	
Honey Tangerines	7	22	47	
Tangelos	14	26	58	
Total	2,211	4,082	4,595	

To subscribe to this report, at no cost, go to the NASS website at http://www.nass.usda.gov/Statistics by State/Florida/Subscribe to FL Reports/index.asp. Complete the Subscribe to FL Reports form, select Florida Crop-Weather and enter your first and last name and your e-mail. The precipitation map used in this report is from the Southeast Regional Climate Center (SERCC) website at http://www.sercc.com/climateinfo/precip_maps. The drought monitor map used in this report is from the U.S. Drought Monitor website at http://droughtmonitor.unl.edu maintained by the National Drought Mitigation Center. The precipitation and temperature data is from the Florida Automated Weather Network (FAWN) at http://fawn.ifas.ufl.edu maintained by UF/IFAS Information Technologies.